

— ABLYBIO, Help Your Research



# Smad7 Rabbit pAb

货号: **AYP25395**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	23kDa/46kDa
实测分子量	46kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 220-390 of human Smad7 (NP_005895.1).
------	---------------------------------------------------------------------------------------------------------------------

## 靶点信息

研究背景	The protein encoded by this gene is a nuclear protein that binds the E3 ubiquitin ligase SMURF2. Upon binding, this complex translocates to the cytoplasm, where it interacts with TGF-beta receptor type-1 (TGFBR1), leading to the degradation of both the encoded protein and TGFBR1. Expression of this gene is induced by TGFBR1. Variations in this gene are a cause of susceptibility to colorectal cancer type 3 (CRCS3). Several transcript variants encoding different isoforms have been found for this gene.
基因ID	4092
基因名	SMAD7
Swiss	O15105 ( <a href="https://www.uniprot.org/uniprotkb/O15105/entry">https://www.uniprot.org/uniprotkb/O15105/entry</a> )
别名	SMAD7,CRCS3,MADH7,MADH8,Smad7 Rabbit pAb,Mothers against decapentaplegic homolog 7,Mothers against decapentaplegic homolog 8

## 产品验证



Western blot analysis of Smad7 expressed in SW480 using Smad7 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)