

— ABLYBIO, Help Your Research



# hnRNP E1/PCBP1 Rabbit pAb

货号: **AYP25415**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>Co-IP:</b> Homo sapiens,Mus musculus <b>WB:</b> Mus musculus
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	37kDa
实测分子量	40KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3,HepG2
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human hnRNP E1/hnRNP E1/PCBP1 (NP_006187.2).
------	--

## 靶点信息

研究背景	This intronless gene is thought to have been generated by retrotransposition of a fully processed PCBP-2 mRNA. This gene and PCBP-2 have paralogues (PCBP3 and PCBP4) which are thought to have arisen as a result of duplication events of entire genes. The protein encoded by this gene appears to be multifunctional. It along with PCBP-2 and hnRNPK corresponds to the major cellular poly(rC)-binding protein. It contains three K-homologous (KH) domains which may be involved in RNA binding. This encoded protein together with PCBP-2 also functions as translational coactivators of poliovirus RNA via a sequence-specific interaction with stem-loop IV of the IRES and promote poliovirus RNA replication by binding to its 5'-terminal cloverleaf structure. It has also been implicated in translational control of the 15-lipoxygenase mRNA, human Papillomavirus type 16 L2 mRNA, and hepatitis A virus RNA. The encoded protein is also suggested to play a part in formation of a sequence-specific alpha-globin mRNP complex which is associated with alpha-globin mRNA stability.
基因ID	5093
基因名	PCBP1
Swiss	Q15365 ( <a href="https://www.uniprot.org/uniprotkb/Q15365/entry">https://www.uniprot.org/uniprotkb/Q15365/entry</a> )
别名	PCBP1,HEL-S-85,HNRPE1,HNRPX,hnRNP-E1,hnRNP-X,hnRNP E1/PCBP1 Rabbit pAb,Alpha-CP1,Heterogeneous nuclear ribonucleoprotein E1,Nucleic acid-binding protein SUB2.3

## 产品验证



Western blot analysis of hnRNP E1/PCBP1 expressed in NIH/3T3, HepG2 using hnRNP E1/PCBP1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)