

— ABLYBIO, Help Your Research



Phospho-IkBa-Y42 Rabbit pAb

货号: **AYP25430**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Rat brain
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	35kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A phospho specific peptide corresponding to residues surrounding Y42 of human IkBa
------	--

靶点信息

研究背景	This gene encodes a member of the NF-kappa-B inhibitor family, which contain multiple ankrin repeat domains. The encoded protein interacts with REL dimers to inhibit NF-kappa-B/REL complexes which are involved in inflammatory responses. The encoded protein moves between the cytoplasm and the nucleus via a nuclear localization signal and CRM1-mediated nuclear export. Mutations in this gene have been found in ectodermal dysplasia anhidrotic with T-cell immunodeficiency autosomal dominant disease.
基因ID	4792
基因名	NFKBIA
Swiss	P25963 (https://www.uniprot.org/uniprotkb/P25963/entry)
别名	IKBA,MAD-3,NFKBI,I κ B alpha,NFKBIA,EDAID2,Phospho-I κ B α -Y42 Rabbit pAb,I-kappa-B-alpha,Major histocompatibility complex enhancer-binding protein MAD3,MAD3

产品验证



Western blot analysis of Phospho-I κ B α -Y42 expressed in HeLa using Phospho-I κ B α -Y42 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)