

AMFR Rabbit pAb

货号: **AYP25433**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	72kDa
实测分子量	72kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3, DU145
细胞定位	Endoplasmic reticulum membrane, Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 384-643 of human AMFR (NP_001135.3).
序列	IADNNRVREEHQGENLDENLVPVAAAEGRPRLNQHNNHFFHFDGSRIASWLPFSFVEMHTTNILGITQASNSQLNAMAHQIQEMFPQVPYHLVLQDLQLTRSVEITTDNILEGRIQVPFPTQRSDSIRPALNSPVERPSSDQEEGETSAQTERVPLDLSPRLETLDFGEVEVEPSEVEDFEARGSRFSKADERQRMLVQRKDELLQARKRFLNKSEDDAASESFLPSEGASSDPVTLRRRMLAAAERRLQKQTS

靶点信息

研究背景	This locus encodes a glycosylated transmembrane receptor. Its ligand, autocrine motility factor, is a tumor motility-stimulating protein secreted by tumor cells. The encoded receptor is also a member of the E3 ubiquitin ligase family of proteins. It catalyzes ubiquitination and endoplasmic reticulum-associated degradation of specific proteins.
------	---

基因ID	267
基因名	AMFR
Swiss	Q9UKV5
别名	AMFR;GP78;RNF45

产品验证



Western blot analysis of AMFR expressed in SKOV3,DU145 using AMFR Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn