

— ABLYBIO, Help Your Research



# Phospho-NF- $\kappa$ B2-S866 Rabbit pAb

货号: **AYP25512**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>IHC:</b> Homo sapiens <b>WB:</b> Mus musculus , Rattus norvegicus , Capra hircus , Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:100 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	47kDa/96kDa
实测分子量	120kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	ovary cancer
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic phosphorylated peptide around S866 of human NF- $\kappa$ B2 (NP_001070962.1).
------	---

## 靶点信息

研究背景	This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFκB). The NFκB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants.
基因ID	4791
基因名	NFKB2
Swiss	Q00653 ( <a href="https://www.uniprot.org/uniprotkb/Q00653/entry">https://www.uniprot.org/uniprotkb/Q00653/entry</a> )
别名	CVID10,H2TF1,LYT-10,LYT10,NF-κB2,p100,p49/p100,p52,Phospho-NF-κB2-S866 Rabbit pAb,NFKB2,DNA-binding factor KBF2,Lymphocyte translocation chromosome 10 protein,Nuclear factor of kappa light polypeptide gene enhancer in B-cells 2,Oncogene Lyt-10

## 产品验证



Western blot analysis of Phospho-NF-κB2-S866 expressed in ovary cancer using Phospho-NF-κB2-S866 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30 μg per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)