

Phospho-NF- κ B2-S870 Rabbit pAb

货号: AYP25516

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	47kDa/96kDa
实测分子量	120kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MDA-MB-435
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	A phospho specific peptide corresponding to residues surrounding S870 of human NF- κ B2
序列	

靶点信息

研究背景	This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NF κ B). The NF κ B complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants.
------	--

基因ID	4791
基因名	NFKB2
Swiss	Q00653
别名	CVID10;H2TF1;LYT-10;LYT10;NF-kB2;p100;p49/p100;p52

产品验证



Western blot analysis of Phospho-NF- κ B2-S870 expressed in MDA-MB-435 using Phospho-NF- κ B2-S870 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn