— ABLYBIO, Help Your Research

PLCG1 (Phospho-Tyr771) Antibody

货号: AYP4048

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC ELISA
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200
理论分子量	148kDa
实测分子量	149kD
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,Jurkat,HepG2,Mouse brain,Mouse lung,Rat brain
细胞定位	Cell projection,lamellipodium,ruffle
纯化	Affinity purification


抗原信息

抗原信息	Synthesized peptide derived from Human PLCG1 (Phospho-Tyr771).

靶点信息

研究背景	The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for this gene.
基因ID	5335
基因名	PLCG1
Swiss	P19174 (https://www.uniprot.org/uniprotkb/P19174/entry)
别名	PLCG1,NCKAP3,PLC-II,PLC1,PLC148,PLCgamma1,PLCG1 (Phospho-Tyr771) Antibody,PLC-148,Phosphoinositide phospholipase C-gamma-1,Phospholipase C-II,Phospholipase C-gamma-1,PLCG1 (Phospho-Tyr771)

产品验证



Western blot detection of PLCG1 (Phospho-Tyr771) in RAW264.7 cell lysates using PLCG1 (Phospho-Tyr771) antibody(1:1000 diluted).

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)