— ABLYBIO, Help Your Research

Androgen Receptor (Phospho-Ser83) Antibody

货号: AYP4127

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC ELISA
推荐浓度	**WB:** 1:500 - 1:2000 **IF/ICC:** 1:50 - 1:200
理论分子量	44kDa/67kDa/99kDa
实测分子量	
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Mouse kidney
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Synthesized peptide derived from Human Androgen Receptor (Phospho-Ser83).

靶点信息

研究背景	The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expansion of the polyglutamine tract from the normal 9-34 repeats to the pathogenic 38-62 repeats causes spinal bulbar muscular atrophy (Kennedy disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Two alternatively spliced variants encoding distinct isoforms have been described.
基因ID	367
基因名	AR
Swiss	P10275 (https://www.uniprot.org/uniprotkb/P10275/entry)
别名	AIS,AR8,DHTR,HUMARA,HYSP1,KD,NR3C4,SBMA,SMAX1,TFM,AR,Androgen Receptor (Phospho-Ser83) Antibody,Dihydrotestosterone receptor,Nuclear receptor subfamily 3 group C member 4,Androgen Receptor (Phospho-Ser83)

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)