

HDAC5 (Phospho-Ser498) Antibody

货号: **AYP4304**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC ELISA
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	112kDa/121kDa/122kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW620,Mouse brain,Mouse heart,Rat brain,Rat heart
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Synthesized peptide derived from Human HDAC5 (Phospho-Ser498).
------	--

靶点信息

研究背景	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events . Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the class II histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. It coimmunoprecipitates only with HDAC3 family member and might form multicomplex proteins. It also interacts with myocyte enhancer factor-2 (MEF2) proteins, resulting in repression of MEF2-dependent genes. This gene is thought to be associated with colon cancer. Two transcript variants encoding different isoforms have been found for this gene.
基因ID	10014

基因名	HDAC5
Swiss	Q9UQL6
别名	HDAC5;HD5;NY-CO-9

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn