

MAP3K8 (Phospho-Ser400) Antibody

货号: **AYP4458**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC ELISA
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	49kDa/52kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	cytoplasm,cytosol
纯化	Affinity purification

抗原信息

抗原信息	Synthesized peptide derived from Human MAP3K8 (Phospho-Ser400).
------	---

靶点信息

研究背景	This gene is an oncogene that encodes a member of the serine/threonine protein kinase family. The encoded protein localizes to the cytoplasm and can activate both the MAP kinase and JNK kinase pathways. This protein was shown to activate I κ B kinases, and thus induce the nuclear production of NF- κ B. This protein was also found to promote the production of TNF- α and IL-2 during T lymphocyte activation. This gene may also utilize a downstream in-frame translation start codon, and thus produce an isoform containing a shorter N-terminus. The shorter isoform has been shown to display weaker transforming activity. Alternate splicing results in multiple transcript variants that encode the same protein.
基因ID	1326

基因名	MAP3K8
Swiss	P41279
别名	MAP3K8;AURA2;COT;EST;ESTF;MEKK8;TPL2;Tpl-2;c-COT

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn