

AKT1/2/3 Antibody

货号: **AYP4798**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC ELISA
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	48kDa/55kDa/51kDa/54kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,Mouse brain,Rat brain
细胞定位	ciliary basal body,cytoplasm,cytosol,microtubule cytoskeleton,mitochondrion,nucleoplasm,nucleus,plasma membrane,spindle
纯化	Affinity purification

抗原信息

抗原信息	Synthesized peptide derived from Human AKT1/2/3.
------	--------------------------------------------------

靶点信息

研究背景	Human AKT serine-threonine protein kinase family includes three members AKT1,AKT2, AKT3, which are also often referred to as protein kinase B alpha, beta, and gamma. These highly similar AKT proteins all have an N-terminal pleckstrin homology domain, a serine/threonine-specific kinase domain and a C-terminal regulatory domain. These proteins are phosphorylated by phosphoinositide 3-kinase (PI3K). AKT/PI3K forms a key component of many signalling pathways that involve the binding of membrane-bound ligands such as receptor tyrosine kinases, G-protein coupled receptors, and integrin-linked kinase. These AKT proteins therefore regulate a wide variety of cellular functions including cell proliferation, survival, metabolism, and angiogenesis in both normal and malignant cells. AKT proteins are recruited to the cell membrane by phosphatidylinositol 3,4,5-trisphosphate (PIP3) after phosphorylation of phosphatidylinositol 4,5-bisphosphate (PIP2) by PI3K. Subsequent phosphorylation of both threonine residue 308 and serine residue 473 is required for full activation of the AKT1 protein encoded by this gene.
基因ID	207,208,10000
基因名	AKT1,AKT2,AKT3
Swiss	P31749/P31751/Q9Y243
别名	AKT1/AKT2/AKT3

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn