

— ABLYBIO, Help Your Research



TRAP220/MED1 Antibody

货号: **AYP5297**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC ELISA
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	61kDa/168kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7
细胞定位	Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Synthesized peptide derived from Human TRAP220/MED1.
------	--

靶点信息

研究背景	The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors. It also regulates p53-dependent apoptosis and it is essential for adipogenesis. This protein is known to have the ability to self-oligomerize.
基因ID	5469
基因名	MED1
Swiss	Q15648 (https://www.uniprot.org/uniprotkb/Q15648/entry)
别名	MED1,CRSP1,CRSP200,DRIP205,DRIP230,PBP,PPARBP,PPARGBP,RB18A,TRAP220,TRIP2,TRAP220/MED1 Antibody,Activator-recruited cofactor 205 kDa component,Mediator complex subunit 1,Peroxisome proliferator-activated receptor-binding protein,Thyroid hormone receptor-associated protein complex 220 kDa component

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)