

MED17 Antibody

货号: **AYP5503**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB ELISA
推荐浓度	WB: 1:500 - 1:2000
理论分子量	15kDa/72kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,A-431,Mouse testis,Rat testis
细胞定位	Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Synthesized peptide derived from Human MED17.
------	---

靶点信息

研究背景	The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors.
基因ID	9440
基因名	MED17

Swiss	Q9NVC6
别名	MED17;CRSP6;CRSP77;DRIP80;TRAP80;SRB4

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn