— ABLYBIO, Help Your Research

# EPHA6 Antibody

**货号: AYP5529**

## 产品信息

| | |
|---|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IHC IF |
| 推荐浓度 | **WB:** 1:500 - 1:2000<br>**IHC:** 1:50 - 1:200<br>**IF:** 1:50 - 1:200 |
| 理论分子量 | 116kD(Calculated). |
| 实测分子量 | 116kD |
| 形式 | Liquid |
| 保存条件 | Store at -20℃. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Brain,Testis |
| 细胞定位 | Membrane ; Single-pass type I |
| 纯化 | Affinity purification |

## 抗原信息


| | |
|---|---|
| 抗原信息 | Synthesized peptide derived from Human EPHA6. |

## 靶点信息

| | |
|---|---|
| 研究背景 | |

| | |
|---|---|
| **基因ID** | 3134 |
| 基因名 | EPHA6 |
| **Swiss** | Q9UF33 (https://www.uniprot.org/uniprotkb/Q9UF33/entry) |
| 别名 | EPHA6 Antibody,EPHA6,EPH homology kinase 2,EPH-like kinase 12,EHK2,HEK12 |

## 产品验证




Western blot analysis of EPHA6 expressed in MCF7 using EPHA6 Antibody at 1:500. Secondary antibody: H RP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120sexpressed in MCF7,SKOV3 using Hes1 Rab bit mAb at 1:500. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s

## 实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)