

— ABLYBIO, Help Your Research



# TAF6L Antibody

货号: **AYP5585**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF ELISA
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF:</b> 1:50 - 1:200
理论分子量	67kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HT-29,Jurkat,Mouse thymus,Rat testis
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息

抗原信息	Synthesized peptide derived from Human TAF6L.
------	---

## 靶点信息

研究背景	Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modifying general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a protein that is a component of the PCAF histone acetylase complex and structurally similar to one of the histone-like TAFs, TAF6. The PCAF histone acetylase complex, which is composed of more than 20 polypeptides some of which are TAFs, is required for myogenic transcription and differentiation.
基因ID	10629
基因名	TAF6L
Swiss	Q9Y6J9 ( <a href="https://www.uniprot.org/uniprotkb/Q9Y6J9/entry">https://www.uniprot.org/uniprotkb/Q9Y6J9/entry</a> )
别名	TAF6L,PAF65A,TAF6L Antibody,PCAF-associated factor 65-alpha

## 产品验证

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))