

PSMD12 Antibody

货号: **AYP5713**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF ELISA
推荐浓度	WB: 1:500 - 1:2000 IF: 1:50 - 1:200
理论分子量	50kDa/52kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,BT-474,HT-29,Mouse heart,Rat heart,Rat lung
细胞定位	cytoplasm,cytosol,extracellular exosome,extracellular region,nucleoplasm
纯化	Affinity purification

抗原信息

抗原信息	Synthesized peptide derived from Human PSMD12.
------	--

靶点信息

研究背景	<p>The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical sub units; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. A pseudogene has been identified on chromosome 3. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.</p>
------	--

基因ID	5718
基因名	PSMD12
Swiss	O00232
别名	PSMD12;Rpn5;p55;STISS

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn