

RAB35 Antibody

货号: **AYP6197**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF ELISA
推荐浓度	WB: 1:500 - 1:2000 IF: 1:50 - 1:200
理论分子量	17kDa/23kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse brain,Rat brain
细胞定位	Cell membrane,Cytoplasmic side,Cytoplasmic vesicle,Endosome,Lipid-anchor,Melanosome,Membrane,clathrin-coated pit,clathrin-coated vesicle
纯化	Affinity purification

抗原信息

抗原信息	Synthesized peptide derived from Human RAB35.
------	---

靶点信息

研究背景	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. That Rab is involved in the process of endocytosis and is an essential rate-limiting regulator of the fast recycling pathway back to the plasma membrane. During cytokinesis, required for the postfurling terminal steps, namely for intercellular bridge stability and abscission, possibly by controlling phosphatidylinositol 4,5-bis phosphate (PIP2 and SEPT2 localization at the intercellular bridge. May indirectly regulate neurite outgrowth. Together with TBC1D13 may be involved in regulation of insulin-induced glucose transporter SLC2A4/GLUT4 translocation to the plasma membrane in adipocytes.
------	--

基因ID	11021
基因名	RAB35
Swiss	Q15286
别名	RAB35;H-ray;RAB1C;RAY

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn