

FGF13 Antibody

货号: **AYP6332**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IF ELISA |
| 推荐浓度 | WB: 1:500 - 1:2000 IF: 1:50 - 1:200 |
| 理论分子量 | 21kDa/22kDa/25kDa/27kDa/28kDa |
| 实测分子量 | |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Mouse brain,Mouse heart,Mouse spinal cord,Rat brain,Rat heart |
| 细胞定位 | Cell projection,Cytoplasm,Cytoplasm,Nucleus,dendrite,filopodium,growth cone,nucleolus |
| 纯化 | Affinity purification |

抗原信息

| | |
|------|---|
| 抗原信息 | Synthesized peptide derived from Human FGF13. |
|------|---|

靶点信息

| | |
|------|--|
| 研究背景 | The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth, and invasion. This gene is located in a region on chromosome X, which is associated with Borjeson-Forssman-Lehmann syndrome (BFLS), making it a possible candidate gene for familial cases of the BFLS, and for other syndromal and nonspecific forms of X-linked mental retardation mapping to this region. Alternative splicing of this gene at the 5' end results in several transcript variants encoding different isoforms with different N-termini. |
| 基因ID | 2258 |

| | |
|-------|------------------------------|
| 基因名 | FGF13 |
| Swiss | Q92913 |
| 别名 | FGF13;FGF-13;FGF2;FHF-2;FHF2 |

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn