

TRAF4 Antibody

货号: **AYP6353**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB ELISA |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 22kDa/53kDa |
| 实测分子量 | |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | MCF7,Mouse liver |
| 细胞定位 | Cell junction,Cell membrane,Cytoplasm,Cytoplasmic side,Nucleus,Peripheral membrane protein,cytoskeleton,perinuclear region,tight junction |
| 纯化 | Affinity purification |

抗原信息

| | |
|------|---|
| 抗原信息 | Synthesized peptide derived from Human TRAF4. |
|------|---|

靶点信息

| | |
|------|--|
| 研究背景 | This gene encodes a member of the TNF receptor associated factor (TRAF) family. TRAF proteins are associated with, and mediate signal transduction from members of the TNF receptor superfamily. The encoded protein has been shown to interact with neurotrophin receptor, p75 (NTR/NTSR1), and negatively regulate NTR induced cell death and NF-kappa B activation. This protein has been found to bind to p47phox, a cytosolic regulatory factor included in a multi-protein complex known as NAD(P)H oxidase. This protein thus, is thought to be involved in the oxidative activation of MAPK8/JNK. Alternatively spliced transcript variants have been observed but the full-length nature of only one has been determined. |
| 基因ID | 9618 |

| | |
|-------|-------------------------|
| 基因名 | TRAF4 |
| Swiss | Q9BUZ4 |
| 别名 | TRAF4;CART1;MLN62;RNF83 |

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn