

— ABLYBIO, Help Your Research



FTCD Rabbit pAb

货号: **AYT1055**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC ICC/IF
推荐浓度	WB 1:500-1:2000, IHC 1:50-1:200: IF :1:50-1:200:
理论分子量	59 kDa
实测分子量	53-62 kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cytoplasm , cytosol .
纯化	Affinity purification

抗原信息


抗原信息	Synthesized peptide derived from human FTCD
------	---

靶点信息

研究背景	
基因ID	10841;

基因名	FTCD
Swiss	O95954 (https://www.uniprot.org/uniprotkb/O95954/entry)
别名	Formimidoyltetrahydrofolate cyclodeaminase; Formimidoyltransferase cyclodeaminase; Formiminotetrahydrofolate cyclodeaminase; Formiminotransferase cyclodeaminase; Formiminotransferase-cyclodeaminase; FTCD; FTCD_HUMAN; Glutamate formiminotransferase; Glutamate formyltransferase; LCHC 1; LCHC1; 58K Golgi protein;

产品验证



The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab FTCD for 1 hour at 37°C. An AlexaFluor 594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green). The nuclear counter stain is DAPI.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)