— ABLYBIO, Help Your Research

IL1RN Rabbit mAb

货号: AYT1160

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC/IF IP
推荐浓度	**WB:** 1:1000 - 1:5000 **IHC 1:100-1:500:** **IF 1:100-1:600:** **IP 1:50-1:200:**
理论分子量	17kDa
实测分子量	20kD
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,PC-3
细胞定位	cytoplasm,extracellular region,extracellular space
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 27-178 of mouse IL1RN (NP_001034790.1).


靶点信息

研究背景	Predicted to enable cytokine receptor binding activity. Acts upstream of or within insulin secretion and lipid metabolic process. Located in extracellular region and vesicle. Human ortholog(s) of this gene implicated in several diseases, including autoimmune disease (multiple); eye disease (multiple); kidney failure (multiple); lung disease (multiple); and vascular disease (multiple). Orthologous to human IL1RN (interleukin 1 receptor antagonist).
基因ID	16181
基因名	Il1rn
Swiss	P25085 (https://www.uniprot.org/uniprotkb/P25085/entry)
别名	IL-1ra,F630041P17Rik,IL1RN Rabbit pAb,Il1rn,IL1 inhibitor

产品验证



Western blot analysis of IL1RN expressed in Hela，K562， RAW264.7 cell and Rat spleen using IL1RN Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Human esophagus was stained with anti-IL1RA (PT1741R) Rabbit antibody

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)