

# MAP2K4 Rabbit pAb

货号: B10289

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens , Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100
理论分子量	44kDa/45kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,BT-474,MCF-7,mouse thymus,mouse liver,rat liver
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 300-399 of human MAP 2K4 (NP_003001.1).	
序列	LATGRFPYPKWNSVFDQLTQVVKGDPPQLSNSEEREFSPSFINFVNLCLTKDESKRPKYKELLKHPFILMYEERAVEVACYV CKILDQMPATPSSPMYVD	

靶点信息

研究背景	This gene encodes a member of the mitogen-activated protein kinase (MAPK) family. Members of this fam ily act as an integration point for multiple biochemical signals and are involved in a wide variety of cellula r processes such as proliferation, differentiation, transcription regulation, and development. They form a three-tiered signaling module composed of MAPKKKS, MAPKKS, and MAPKS. This protein is phosphorylated at serine and threonine residues by MAPKKKS and subsequently phosphorylates downstream MAPK target s at threonine and tyrosine residues. A similar protein in mouse has been reported to play a role in liver o rganogenesis. A pseudogene of this gene is located on the long arm of chromosome X. Alternative splicin g results in multiple transcript variants.
基因 <b>ID</b>	6416
基因名	MAP2K4
Swiss	P45985
别名	MAP2K4;JNKK;JNKK1;MAPKK4;MEK4;MKK4;PRKMK4;SAPKK-1;SAPKK1;SEK1;SERK1;SKK1

## 产品验证



Western blot analysis of extracts of various cell lines, using MAP2K4 antibody at 1:1000 dilution.S econdary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.Lysates/proteins: 25ug pe r lane.Blocking buffer: 3% nonfat dry milk in TBST.Detection: ECL Basic Kit.Exposure time: 90s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn