

# B4GALT4 Rabbit pAb

货号: B10744

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>IHC:</b> Human breast cancer <b>WB:</b> Human breast cancer
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	40kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HT-29,Mouse kidney,Mouse thymus,Mouse lung,Rat kidney,Rat thymus,Rat lung
细胞定位	Golgi apparatus,Golgi stack membrane,Single-pass type II membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 35-344 of human B4GAL T4 (NP_003769.1).
序列	VGAIQEIPKAKEFMANFHKTLLGKGKTLTNEASTKKVELDNCPSVSPYLRGQSKLIFKPDLTLEEVQAENPKVSRGRYRPQE CKALQRVAILVPHRNREKHLMLLEHLHPFLQRQQLDYGIYVIHQAEKKFNRAKLLNVGYLEALKENWDCFIFHDVDLVP ENDFNLYKCEEHPKHLVVGRNSTGYRLRSGYFGGVVTALSREQFFKVNGFSNNYWGWGGEDDDLRLRVELQRMKISRP LPEVGKYTMVFHTRDKGNEVNAERMKLLHQVSRVWRTDGLSSCSYKLVSVEHNPLYINITVDFWFGA

## 靶点信息

研究背景	This gene is one of seven beta-1,4-galactosyltransferase (beta4GalT) genes. They encode type II membrane-bound glycoproteins that appear to have exclusive specificity for the donor substrate UDP-galactose; a II transfer galactose in a beta1,4 linkage to similar acceptor sugars: GlcNAc, Glc, and Xyl. Each beta4GalT has a distinct function in the biosynthesis of different glycoconjugates and saccharide structures. As type II membrane proteins, they have an N-terminal hydrophobic signal sequence that directs the protein to the Golgi apparatus and which then remains uncleaved to function as a transmembrane anchor. By sequence similarity, the beta4GalTs form four groups: beta4GalT1 and beta4GalT2, beta4GalT3 and beta4GalT4, beta4GalT5 and beta4GalT6, and beta4GalT7. The enzyme encoded by this gene appears to mainly play a role in glycolipid biosynthesis. Two alternatively spliced transcript variants have been found for this gene.
基因ID	8702
基因名	B4GALT4
Swiss	O60513
别名	B4GALT4;B4Gal-T4;beta4Gal-T4;beta-1

## 产品验证



Western blot analysis of B4GALT4 expressed in Mouse kidney, Rat kidney, 293T using B4GALT4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 S.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)