

KIR3DL3 Rabbit pAb

货号: **B10753**

产品信息

反应 Human 宿主 Rabbit 克隆性 Polyclonal 预测反应 应用 WB IHC 推荐浓度 WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
克隆性 Polyclonal 预测反应 应用 WB IHC 推荐浓度 WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
应用 WB IHC 推荐浓度 WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
推荐浓度 WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
IHC: 1:50 - 1:200
理论分子量 44kDa
实测分子量 45kDa
形式 Liquid
保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物 Unconjugated
阳性对照 HepG2,SGC-7901
细胞定位 Cell membrane,Single-pass type I membrane protein
纯化 Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 26-322 of human KIR3D L3 (NP_703144.3).
序列	QDKPFLSAWPGTVVSEGQHVTLQCRSRLGFNEFSLSKEDGMPVPELYNRIFRNSFLMGPVTPAHAGTYRCCSSHPHSPT GWSAPSNPVVIMVTGVHRKPSLLAHPGPLVKSGETVILQCWSDVRFERFLLHREGITEDPLRLVGQLHDAGSQVNYSMGP MTPALAGTYRCFGSVTHLPYELSAPSDPLDIVVVGLYGKPSLSAQPGPTVQAGENVTLSCSSRSLFDIYHLSREAEAGELRL TAVLRVNGTFQANFPLGPVTHGGNYRCFGSFRALPHAWSDPSDPLPVSVTGNSRNL

靶点信息

研究背景	Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural kil ler cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several 'framework' genes are found in all hapl otypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracell ular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic do main. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain la ck the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response. This gene is one of the 'framework' loci that is present on all haplotypes.
基因 ID	115653
基因名	
Swiss	Q8N743
别名	KIR3DL3;CD158Z;KIR3DL7;KIR44;KIRC1

产品验证



Western blot analysis of KIR3DL3 expressed in HepG2,SGC-7901 using KIR3DL3 Rabbit pAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking b uffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn