

OR2S2 Rabbit pAb

货号: **B10824**

产品信息

反应	Human, Mouse, Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	35kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HepG2
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 200-300 of human OR2S2 (NP_0639 50.2).
序列	MEVTNVIFLGVPVLFISFSYVFIITTILRIPSAEGRKKVFSTCSAHLTVVIVFYGTLFFMYGKPKSKDSMGADKEDLSDKLIPLF YGVVTPMLNPIIYSLR

靶点信息

研究背景	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that trigg ers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-c oupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmemb rane domain structure with many neurotransmitter and hormone receptors and are responsible for the re cognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional olfactory receptor, while other individuals have an allele encoding a protein that is predicted to be non-functional.
基因 ID	56656
基因名	OR2S2
Swiss	Q9NQN1
别名	OR2S2;OR37A;OST715

产品验证



Western blot analysis of OR2S2 expressed in U-87MG,HepG2 using OR2S2 Rabbit pAb at 1:1000. Seconda ry antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn