

# **APEX2** Rabbit pAb

货号: **B10843** 

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	57kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,A-549
细胞定位	Cytoplasm,Mitochondrion,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-180 of human APEX2 ( NP_055296.2).
序列	MLRVVSWNINGIRRPLQGVANQEPSNCAAVAVGRILDELDADIVCLQETKVTRDALTEPLAIVEGYNSYFSFSRNRSGYSG VATFCKDNATPVAAEEGLSGLFATQNGDVGCYGNMDEFTQEELRALDSEGRALLTQHKIRTWEGKEKTLTLINVYCPHAD PGRPERLVFKMRFYRLLQI

靶点信息

研究背景	Apurinic/apyrimidinic (AP) sites occur frequently in DNA molecules by spontaneous hydrolysis, by DNA da maging agents or by DNA glycosylases that remove specific abnormal bases. AP sites are pre-mutagenic I esions that can prevent normal DNA replication so the cell contains systems to identify and repair such si tes. Class II AP endonucleases cleave the phosphodiester backbone 5' to the AP site. This gene encodes a protein shown to have a weak class II AP endonuclease activity. Most of the encoded protein is located in the nucleus but some is also present in mitochondria. This protein may play an important role in both nuc lear and mitochondrial base excision repair. Alternatively spliced transcript variants encoding multiple iso forms have been observed for this gene.
基因ID	27301
基因名	APEX2
Swiss	Q9UBZ4
别名	APEX2;APE2;APEXL2;XTH2;ZGRF2

## 产品验证



Western blot analysis of APEX2 expressed in U-87MG,A-549 using APEX2 Rabbit pAb at 1:1000. Secondar y antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn