

RFX1 Rabbit pAb

货号: **B10867**

产品信息

| 反应 | Human |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 104kDa |
| 实测分子量 | 130kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | LO2 |
| 细胞定位 | Nucleus |
| 纯化 | Affinity purification |

抗原信息


| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 200-300 of human RFX1 (NP_002909 .4). |
|------|---|
| 序列 | QQVHSPPEQSPVQANSSSSKTAGAPTGTVPQQLQVHGVQQSVPVTQERSVVQATPQAPKPGPVQPLTVQGLQPVHVA QEVQQLQQVPVPHVYSSQVQYVEG |

靶点信息

| 研究背景 | This gene encodes a member of the regulatory factor X (RFX) family of transcription factors, which are ch aracterized by a winged-helix DNA-binding domain. The encoded transcription factor contains an N-termi nal activation domain and a C-terminal repression domain, and may activate or repress target gene expression depending on cellular context. This transcription factor has been shown to regulate a wide variety of genes involved in immunity and cancer, including the MHC class II genes and genes that may be involved in cancer progression. This gene exhibits altered expression in glioblastoma and the autoimmune dise ase systemic lupus erythematosis (SLE). |
|------|---|

| 基因ID | 5989 |
|-------|--------------|
| 基因名 | RFX1 |
| Swiss | P22670 |
| 别名 | RFX1;EFC;RFX |

产品验证



Western blot analysis of RFX1 expressed in LO2 using RFX1 Rabbit pAb at 1:1000. Secondary antibody: H RP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn