

TriMethyl-Histone H3-K27 Rabbit pAb

货号: B10924

产品信息

反应	Human, Mouse, Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IF: Human embryonic stem cells , Mouse E16.5 neocortex , Goat sperm , HEK293T , Rice , Mus musculus , Gallus gallus , Sus scrofa , Homo sapiens ChIP: A549 , Mouse embryonic stem cells , HeLa , Human , Rice , Gossypium , B. napus , Oryza sativa , Mus musculus , Homo sapiens , Rattus norvegicus , castor bean IHC: Mouse lung tumor , Mus musculus , Homo sapiens , Sus scrofa WB: Mus musculus , Homo sapiens , Drosophila , Rattus norvegicus CHIP: Arabidopsis thaliana , castor bean , Mus musculus , Homo sapiens IP: Gossypium , Arabidopsis , Mus musculus CoIP: N. benthamiana ChiP: Mus musculus
应用	WB IHC IP CHIP
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:50 - 1:200 ChIP: 1:20 - 1:100
理论分子量	15kDa
实测分子量	17KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7 acid extract,293T acid extract,C2C12 acid extract,NIH/3T3 acid extract,C6 acid extract,H3 protein
细胞定位	Chromosome, Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic trimethylated peptide around K27 of human histone H3 (NP_003520.1).
序列	AARKS

靶点信息

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octa mer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent his tone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.
8290
HIST3H3
Q16695
H3.4;H3/g;H3FT;H3t;HIST3H3;Histone H3;HIST1H3A

产品验证



Western blot analysis of TriMethyl-Histone H3-K27 expressed in MCF7 acid extract,293T acid extract, ct,C2C12 acid extract using TriMethyl-Histone H3-K27 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn