

TriMethyl-Histone H3-K4 Rabbit pAb

货号: B10926

产品信息


反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Human cells , HEK293T , Mouse Bone marrow stromal cells , Homo sapiens , Saccharomyces cerevi siae , Mus musculus , Rattus norvegicus , Oryza sativa , Shrimp , Drosophila ChIP: Plant samples , Mouse primary calvarial cells , B. napus , Mus musculus , Oryza sativa , Arabidopsi s thaliana , Homo sapiens , Rattus norvegicus IF: HEK293T , Mouse E16.5 neocortex , Raw264.7 , Mus musculus , Oryza sativa , Rattus norvegicus , Su s scrofa , Homo sapiens CHIP: Arabidopsis thaliana , Zea mays , Homo sapiens , Mus musculus IP: Homo sapiens , Mus musculus IHC: Mus musculus , Rattus norvegicus , Bombyx mori
应用	WB IHC CHIP
推荐浓度	DB: 1:500 - 1:2000 WB: 1:100 - 1:500 IHC: 1:50 - 1:200 ChIP: 1:20 - 1:100
理论分子量	15kDa
实测分子量	17KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3
细胞定位	Chromosome, Nucleus
纯化	Affinity purification

抗原信息

抗原信息	A synthetic trimethylated peptide around K4 of human histone H3 (NP_003520.1).
序列	RTKQT

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octa mer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent his tone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.
基因 ID	8290
基因名	HIST3H3
Swiss	Q16695
别名	H3.4;H3/g;H3FT;H3t;HIST3H3;Histone H3;HIST1H3A

产品验证



Western blot analysis of TriMethyl-Histone H3-K4 expressed in NIH/3T3 using TriMethyl-Histone H3-K4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/protein s: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn