

TriMethyl-Histone H3-K36 Rabbit pAb

货号: B10932

产品信息


| 反应 | Human,Mouse,Rat |
|-------|--|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | IF: HEK293T, Rattus norvegicus, Homo sapiens, Mus musculus WB: HEK293T, Hippocampus, Homo sapiens, Yeast, Saccharomyces cerevisiae, Mus musculus, Rattus norvegicus, Oryza sativa IHC: Rattus norvegicus, Sus scrofa ChIP: Oryza sativa |
| 应用 | WB IHC IP CHIP |
| 推荐浓度 | WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:50 - 1:200 ChIP: 1:20 - 1:100 |
| 理论分子量 | 15kDa |
| 实测分子量 | 17KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | C6,H3 |
| 细胞定位 | Chromosome, Nucleus |
| 纯化 | Affinity purification |

抗原信息

| 抗原信息 | A synthetic trimethylated peptide around K36 of human histone H3 (NP_003520.1). |
|------|---|
| 序列 | GVKKP |

| 研究背景 | Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octa mer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is fu rther compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes t o form higher order chromatin structures. This gene is intronless and encodes a replication-dependent his tone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3. |
|-------|---|
| 基因ID | 8290 |
| 基因名 | HIST3H3 |
| Swiss | Q16695 |
| 别名 | H3.4;H3/g;H3FT;H3t;HIST3H3;Histone H3;HIST1H3A |

产品验证



Western blot analysis of TriMethyl-Histone H3-K36 expressed in C6,H3 using TriMethyl-Histone H3-K36 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysate s/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn