

DiMethyl-Histone H3-K27 Rabbit pAb

货号: B10948

产品信息

反应 Human,Mouse,Rat 宿主 Rabbit 克隆性 Polyclonal
克隆性 Polyclonal
预测反应 WB: 293FT,PC9 , HeLa , Homo sapiens IF: Goat sperm
应用 WB IP CHIP
推荐浓度 WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200 IP: 1:50 - 1:200 ChIP: 1:20 - 1:100
理论分子量 15kDa
实测分子量 17KDa
形式 Liquid
保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物 Unconjugated
阳性对照 HeLa acid extract,NIH/3T3 acid extract,C6 acid extract,H3 protein
细胞定位 Chromosome,Nucleus
纯化 Affinity purification


抗原信息

抗原信息	A synthetic dimethylated peptide around K27 of human histone H3 (NP_003520.1).
序列	ARKSA

靶点信息

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octa mer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent his tone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.
基因ID	8290
基因名	HIST3H3
Swiss	Q16695
别名	H3.4;H3/g;H3FT;H3t;HIST3H3;Histone H3;HIST1H3A

产品验证



Western blot analysis of DiMethyl-Histone H3-K27 expressed in HeLa acid extract,NIH/3T3 acid extract,C6 acid extract using DiMethyl-Histone H3-K27 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn