

# Acetyl-Histone H3-K27 Rabbit pAb

货号: B11012

#### 产品信息


反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Prawn gills , Mus musculus , Homo sapiens , Sus scrofa , Danio rerio , Rattus norvegicus , Monascus ruber ChIP: Mus musculus , B. napus , Oryza sativa , Homo sapiens , Rattus norvegicus , Gallus gallus CHIP: Arabidopsis thaliana , Homo sapiens IP: Rattus norvegicus IF: Homo sapiens , Mus musculus IHC: Rattus norvegicus CoIP: Rattus norvegicus Co-IP: Rattus norvegicus
应用	WB IHC IP CHIP
推荐浓度	WB: 1:1000 - 1:5000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:50 - 1:200 ChIP: 1:20 - 1:100
理论分子量	15kDa
实测分子量	17KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3,C6
细胞定位	Chromosome, Nucleus
纯化	Affinity purification

## 抗原信息

抗原信息	A synthetic acetylated peptide around K27 of human Histone H3 (NP_003520.1).
序列	AARKS

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octa mer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent his tone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.
基因 <b>ID</b>	8290
基因名	HIST3H3
Swiss	P68431
别名	H3.4;H3/g;H3FT;H3t;HIST3H3;Histone H3;HIST1H3A

#### 产品验证



Western blot analysis of Acetyl-Histone H3-K27 expressed in NIH/3T3,C6 using Acetyl-Histone H3-K2 7 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. Lysates/pro teins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exp osure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn