

MyD88 Rabbit pAb

货号: B11040

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: CAL27 , Mus musculus , Rattus norvegicus , Homo sapiens IF: Mus musculus IB: Homo sapiens IHC: Homo sapiens , Mus musculus , Rattus norvegicus
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	15kDa/20kDa/28kDa/31kDa/33kDa/34kDa
实测分子量	33KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,Mouse lung,Mouse uterus,Rat lung
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 50-150 of human MyD88 (NP_001166038.2).
序列	LAEEMDFEYLEIRQLETQADPTGRLLDAWQGRPGASVGRLLLELLTKLGRDDVLLELGPSIEEDCQKYILKQQQEAEKPLQVAAVDSSVPRTAELAGITTL

靶点信息

研究背景	This gene encodes a cytosolic adapter protein that plays a central role in the innate and adaptive immune response. This protein functions as an essential signal transducer in the interleukin-1 and Toll-like receptor signaling pathways. These pathways regulate that activation of numerous proinflammatory genes. The encoded protein consists of an N-terminal death domain and a C-terminal Toll-interleukin1 receptor domain. Patients with defects in this gene have an increased susceptibility to pyogenic bacterial infections. Alternate splicing results in multiple transcript variants.
基因ID	4615
基因名	MYD88
Swiss	Q99836
别名	MYD88;MYD88D;myeloid differentiation primary response 88;MyD88

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn