

CD11b/ITGAM Rabbit pAb

货号: B11089

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | IHC: Mus musculus , Rattus norvegicus , Homo sapiens IF: Homo sapiens , Mus musculus , Rattus norvegicus WB: Mus musculus , Rattus norvegicus FC: Homo sapiens |
| 应用 | WB IHC IF/ICC |
| 推荐浓度 | WB: 1:100 - 1:500 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 127kDa |
| 实测分子量 | 170KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | THP-1,U-937,Mouse lung,Mouse brain,Mouse spleen,Rat lung |
| 细胞定位 | Membrane,Single-pass type I membrane protein |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 145-340 of CD11b/ITGA M (NP_000623.2). |
| 序列 | PQEDSDIAFLIDGSIGSIIPHDFRMKEFVSTVMEQLKKSKTLFSLMQYSEEFRIHFTFKEFQNNPNPRSLVKPITQLLGRTHT ATGIRKVVRLEFNITNGARKNAFKILVVITDGEKGFDPLGYEDVPEADREGVIRYVIGVGDAFRSEKSRSRQELNTIASKPPRDHV FQVNNEFAALKTIQNQLREKIFAIEGTQT |

靶点信息

| | |
|-------|---|
| 研究背景 | This gene encodes the integrin alpha M chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form a leukocyte-specific integrin referred to as macrophage receptor 1 ('Mac-1'), or inactivated-C3b (iC3b) receptor 3 ('CR3'). The alpha M beta 2 integrin is important in the adherence of neutrophils and monocytes to stimulated endothelium, and also in the phagocytosis of complement coated particles. Multiple transcript variants encoding different isoforms have been found for this gene. |
| 基因ID | 3684 |
| 基因名 | ITGAM |
| Swiss | P11215 |
| 别名 | CD11B;CR3A;MAC-1;MAC1A;MO1A;SLEB6;ITGAM;CD11b/ITAM |

产品验证



Western blot analysis of CD11b/ITGAM expressed in Mouse lung, Rat lung, THP-1 using CD11b/ITGAM Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn