

TGF beta 1 Rabbit pAb

货号: B11115

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IF: Homo sapiens , Mus musculus WB: Mus musculus , Homo sapiens , Rattus norvegicus IHC: Mus musculus , Homo sapiens
应用	IF/ICC
推荐浓度	IF/ICC: 1:50 - 1:200
理论分子量	44kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Secreted,extracellular matrix,extracellular space
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 30-278 of human TGF b eta 1 (NP_000651.3).
序列	LSTCKTIDMELVKRKRIEAI RGQILSKRLASPPSQGEVPPGPLPEAVLALYNSTRDRVAGESAEPEPEPEADYYAKEVTRVLM VETHNEIYDKFKQSTHSIYMFFNTSELREAVPEPVLLSRAELRLRLKLKVEQHVELYQKYSNNSWRYLSNRLLAPSDSPEW LSFDTVGVVRQWLSRGGEIEGFRLSAHCSDSRDNTLQVDINGFTTGRRGDLATIHGMNRPFLLLMATPLERAQHLQSSR HRR

靶点信息

研究背景	This gene encodes a member of the transforming growth factor beta (TGFB) family of cytokines, which are multifunctional peptides that regulate proliferation, differentiation, adhesion, migration, and other functions in many cell types. Many cells have TGFB receptors, and the protein positively and negatively regulates many other growth factors. The secreted protein is cleaved into a latency-associated peptide (LAP) and a mature TGFB1 peptide, and is found in either a latent form composed of a TGFB1 homodimer, a LAP homodimer, and a latent TGFB1-binding protein, or in an active form composed of a TGFB1 homodimer. The mature peptide may also form heterodimers with other TGFB family members. This gene is frequently upregulated in tumor cells, and mutations in this gene result in Camurati-Engelmann disease.
基因ID	7040
基因名	TGFB1
Swiss	P01137
别名	TGFB1;CED;DPD1;LAP;TGFB;TGFBeta;TGF-β

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn