

Rad50 Rabbit pAb

货号: B11256

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mouse adipose-derived stem cells , HCC827 , Mus musculus , Homo sapiens IF: Mus musculus IHC: Rattus norvegicus
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	138kDa/153kDa/154kDa
实测分子量	154KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,K-562,U-937,Mouse testis,Mouse thymus,Rat testis
细胞定位	Chromosome,Nucleus,telomere
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Rad50 (NP_005723.2).
序列	MSRIEKMSILGVRSFGIEDKDKQIITFFSPLTILVGPNAGKTTIIECLKYICTGDFPPGKGNTFVHDPKVAQETDVRAQIRLQFRDVNGELIAVQRSM

靶点信息

研究背景	The protein encoded by this gene is highly similar to <i>Saccharomyces cerevisiae</i> Rad50, a protein involved in DNA double-strand break repair. This protein forms a complex with MRE11 and NBS1. The protein complex binds to DNA and displays numerous enzymatic activities that are required for nonhomologous joining of DNA ends. This protein, cooperating with its partners, is important for DNA double-strand break repair, cell cycle checkpoint activation, telomere maintenance, and meiotic recombination. Knockout studies of the mouse homolog suggest this gene is essential for cell growth and viability. Mutations in this gene are the cause of Nijmegen breakage syndrome-like disorder.
基因ID	10111
基因名	RAD50
Swiss	Q92878
别名	RAD50;NBSLD;RAD502;hRad50

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn