

Goat Anti-Mouse IgG (H+L)

货号: B11336

产品信息

反应	
宿主	Goat
克隆性	
预测反应	Co-IP: Homo sapiens WB: Mus musculus,Homo sapiens , Cavia porcellus FC: Mus musculus
应用	
推荐浓度	
理论分子量	
实测分子量	25KDa/55KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse IgG
细胞定位	
纯化	Affinity purification

抗原信息

抗原信息	Mouse IgG
序列	

靶点信息

研究背景	Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies . Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.
------	---

基因ID	
基因名	
Swiss	
别名	

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn