

# TIMP2 Rabbit pAb

货号: **B11355** 

### 产品信息

反应 Human,Mouse,Rat 宿主 Rabbit	
宿主 Rabbit	
克隆性 Polyclonal	
预测反应  WB: MDA-MB231 , Homo sapiens  IF: Homo sapiens,Mus musculus  IHC: Mus musculus	
应用 WB IHC IF/ICC	
推荐浓度  WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200	
理论分子量 24kDa	
实测分子量 24KDa	
形式 Liquid	
保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.	
偶联物 Unconjugated	
阳性对照 A-431,Mouse testis,Rat testis	
细胞定位 Secreted	
纯化 Affinity purification	


### 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 50-150 of human TIMP2 (NP_003246. 1).
序列	VSEKEVDSGNDIYGNPIKRIQYEIKQIKMFKGPEKDIEFIYTAPSSAVCGVSLDVGGKKEYLIAGKAEGDGKMHITLCDFIVP WDTLSTTQKKSLNHRYQM

靶点信息

研究背景	This gene is a member of the TIMP gene family. The proteins encoded by this gene family are natural inhi bitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extracellular matrix. In addition to an inhibitory role against metalloproteinases, the encoded protein has a unique role among TIMP family members in its ability to directly suppress the proliferation of endothelial cells. As a re sult, the encoded protein may be critical to the maintenance of tissue homeostasis by suppressing the proliferation of quiescent tissues in response to angiogenic factors, and by inhibiting protease activity in tis sues undergoing remodelling of the extracellular matrix.
基因ID	7077
基因名	TIMP2
Swiss	P16035
别名	TIMP2;CSC-21K;DDC8

### 产品验证



Western blot analysis of TIMP2 expressed in A-431,Mouse testis,Rat testis using TIMP2 Rabbit pAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn