

TGFB2 Rabbit pAb

货号: **B11375**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens , Mus musculus , Gallus gallus IF: Gallus gallus
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	47kDa/50kDa
实测分子量	48KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung, Mouse heart, Mouse uterus, Rat lung, Rat heart
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 60-220 of human TGFB2 (NP_003229.1).	
序列	PEEVPPEVISIYNSTRDLLQEKASRRAAACERERSDEEYYAKEVYKIDMPPFFPSENAIPPTFYRPYFRIVRFDVSAMEKNASN LVKAEFRVFRLQNPKARVPEQRIELYQILKSKDLTSPTQRYIDSKVVKTRAEGEWLSFDVTDAVHEWLHHKDRNLGF	

靶点信息

研究背景	This gene encodes a member of the transforming growth factor beta (TGFB) family of cytokines, which ar e multifunctional peptides that regulate proliferation, differentiation, adhesion, migration, and other funct ions in many cell types by transducing their signal through combinations of transmembrane type I and ty pe II receptors (TGFBR1 and TGFBR2) and their downstream effectors, the SMAD proteins. Disruption of t he TGFB/SMAD pathway has been implicated in a variety of human cancers. The encoded protein is secre ted and has suppressive effects of interleukin-2 dependent T-cell growth. Translocation t(1;7)(q41;p21) b etween this gene and HDAC9 is associated with Peters' anomaly, a congenital defect of the anterior cham ber of the eye. The knockout mice lacking this gene show perinatal mortality and a wide range of develop mental, including cardiac, defects. Alternatively spliced transcript variants encoding different isoforms ha ve been identified.
基因 ID	7042
基因名	TGFB2
Swiss	P61812
别名	TGFB2;G-TSF;LDS4;TGF-beta2

产品验证



Western blot analysis of TGFB2 expressed in Mouse lung,Rat lung,Mouse heart using TGFB2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per I ane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn