

Neutrophil Elastase (ELANE) Rabbit pAb

货号: B11388

产品信息


反应 Human,Mouse,Rat	
宿主 Rabbit	
克隆性 Polyclonal	
预测反应 WB: Homo sapiens , Rattus norvegicus IF: Mus musculus , Homo sapiens	
应用 WB IHC IF/ICC	
推荐浓度 WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200	
理论分子量 28kDa	
实测分子量 30KDa	
形式 Liquid	
保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.	
偶联物 Unconjugated	
阳性对照 Rat spleen	
细胞定位 cell surface,cytoplasm,cytosol,extracellular exosome,extracellular region,extracellular space,phagesicle	gocytic v
纯化 Affinity purification	

抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 168-267 of human Neutrophil Elasta se (ELANE) (NP_001963.1).
序列	VLQELNVTVVTSLCRRSNVCTLVRGRQAGVCFGDSGSPLVCNGLIHGIASFVRGGCASGLYPDAFAPVAQFVNWIDSIIQR SEDNPCPHPRDPDPASRTH

研究背景	Elastases form a subfamily of serine proteases that hydrolyze many proteins in addition to elastin. Huma ns have six elastase genes which encode structurally similar proteins. The encoded preproprotein is prote olytically processed to generate the active protease. Following activation, this protease hydrolyzes proteins within specialized neutrophil lysosomes, called azurophil granules, as well as proteins of the extracellu lar matrix. The enzyme may play a role in degenerative and inflammatory diseases through proteolysis of collagen-IV and elastin. This protein also degrades the outer membrane protein A (OmpA) of E. coli as well as the virulence factors of such bacteria as Shigella, Salmonella and Yersinia. Mutations in this gene are associated with cyclic neutropenia and severe congenital neutropenia (SCN). This gene is present in a gene cluster on chromosome 19.
基因 ID	1991
基因名	ELANE
Swiss	P08246
别名	ELANE;ELA2;GE;HLE;HNE;NE;PMN-E;SCN1

产品验证



Western blot analysis of Neutrophil Elastase (ELANE) expressed in Rat spleen using Neutrophil Elastas e (ELANE) Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lys ates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn