

STAT6 Rabbit pAb

货号: **B11482**

产品信息

Human,Mouse,Rat
Rabbit
Polyclonal
WB: Mus musculus , Rattus norvegicus , Homo sapiens
WB IF/ICC
WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
74kDa/81kDa/94kDa
110KDa
Liquid
Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
Unconjugated
HeLa,Jurkat,PC-12,NIH/3T3,Mouse lung,Mouse spleen,Rat lung
Cytoplasm,Nucleus
Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 550-650 of human STAT6 (NP_00314 4.3).
序列	LLLNEPDGTFLLRFSDSEIGGITIAHVIRGQDGSPQIENIQPFSAKDLSIRSLGDRIRDLAQLKNLYPKKPKDEAFRSHYKPEQ MGKDGRGYVPATIKMTV

靶点信息

研究背景	The protein encoded by this gene is a member of the STAT family of transcription factors. In response to c ytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinase s, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcripti on activators. This protein plays a central role in exerting IL4 mediated biological responses. It is found to induce the expression of BCL2L1/BCL-X(L), which is responsible for the anti-apoptotic activity of IL4. Knoc kout studies in mice suggested the roles of this gene in differentiation of T helper 2 (Th2) cells, expressio n of cell surface markers, and class switch of immunoglobulins. Alternative splicing results in multiple transcript variants.
基因ID	6778
基因名	STAT6
Swiss	P42226
别名	D12S1644;IL-4-STAT;STAT6B;STAT6C;STAT6

产品验证



Western blot analysis of STAT6 expressed in Mouse lung,Rat lung,HeLa using STAT6 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn