

SIRT6 Rabbit pAb

货号: B11539

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	36kDa/39kDa
实测分子量	39-45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,Mouse spleen,Mouse stomach,A-549,LO2,BT-474,HL-60,Mouse heart
细胞定位	Nucleus,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 200 to the C-terminus of mouse SIRT6 (NP_001156902.1).
序列	NLQPTKHDRQADLRIHGYVDEVMCRLMKHLGLEIPAWDGPCVLDKALPPLPRPVALKAEPVVHNGAVHVSYKSKPNSPIL HRPPKRVKTEAAPS

靶点信息

研究背景	This gene encodes a member of the sirtuin family of NAD-dependent enzymes that are implicated in cellular stress resistance, genomic stability, aging and energy homeostasis. The encoded protein is localized to the nucleus, exhibits ADP-ribosyl transferase and histone deacetylase activities, and plays a role in DNA repair, maintenance of telomeric chromatin, inflammation, lipid and glucose metabolism. Alternative splicing results in multiple transcript variants encoding different isoforms.
基因ID	50721

基因名	
Swiss	Q8N6T7
别名	SIRT6;SIR2L6;sirtuin 6

产品验证



Western blot analysis of SIRT6 expressed in Mouse spleen, HeLa, 293T using SIRT6 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn