

# K48-linkage Specific Polyubiquitin Rabbit pAb

货号: B11616

#### 产品信息

反应	Human,Mouse,Rat,Other (Wide Range)
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Drosophila melanogaster , Homo sapiens , Monkey
应用	WB DB
推荐浓度	<b>WB:</b> 1:100 - 1:500 <b>DB:</b> 1:500 - 1:1000
理论分子量	
实测分子量	15-250KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,HeLa,C2C12,C6,PC-12
细胞定位	
纯化	Affinity purification


#### 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence containing polyubiquitinated protein (K48 linkage).
序列	

靶点信息

研究背景	Protein ubiquitination is a dynamic multifaceted post-translational modification involved in nearly all aspects of eukaryotic biology. Owing to its molecular features, ubiquitin can form various homo- and heteroty pic polymers on substrate proteins, thereby provoking distinct cellular responses. Ubiquitin can be ubiquitinated on seven lysine (Lys) residues or on the N-terminus, leading to polyubiquitin chains that can encompass complex topologies. Polyubiquitin chains have different functions depending on the Lys residue of the ubiquitin that is linked: Lys-6-linked may be involved in DNA repair; Lys-11-linked is involved in ERAD (endoplasmic reticulum-associated degradation) and in cell-cycle regulation; Lys-29-linked is involved in lysosomal degradation; Lys-33-linked is involved in kinase modification; Lys-48-linked is involved in protein degradation via the proteasome; Lys-63-linked is involved in endocytosis, DNA-damage responses as well as in signaling processes leading to activation of the transcription factor NF-kappa-B.
基因 <b>ID</b>	
基因名	
Swiss	
别名	

## 产品验证



Western blot analysis of K48-linkage Specific Polyubiquitin expressed in Jurkat,HeLa,C2C12 usin g K48-linkage Specific Polyubiquitin Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry m ilk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn