

# YTHDF1 Rabbit pAb

货号: B11617

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>IHC:</b> Homo sapiens <b>WB:</b> Homo sapiens
应用	<a href="#">WB</a> <a href="#">IHC</a>
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IHC:</b> 1:50 - 1:200
理论分子量	20kDa/60kDa
实测分子量	70KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,Mouse testis,Rat thymus
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 280-390 of human YTHDF1 (NP_060268.2).
序列	VPKAPVPQQAPSPQAAPQPQQVAQLPAQPPALAQPQYQSPQPPQTRWVAPRNRNAAFGQSGGAGSDSNSPGNVQPNSAPSVESHVPLEKLKAAHSYNPKFEWNLKSGR

靶点信息

研究背景	Specifically recognizes and binds N6-methyladenosine (m6A-containing mRNAs, and regulates their stability. M6A is a modification present at internal sites of mRNAs and some non-coding RNAs and plays a role in mRNA stability and processing. Acts as a regulator of mRNA stability by promoting degradation of m6A-containing mRNAs via interaction with the CCR4-NOT complex. The YTHDF paralogs (YTHDF1, YTHDF2 and YTHDF3) shares m6A-containing mRNAs targets and act redundantly to mediate mRNA degradation and cellular differentiation. Required to facilitate learning and memory formation in the hippocampus by binding to m6A-containing neuronal mRNAs (By similarity. Acts as a regulator of axon guidance by binding to m6A-containing ROBO3 transcripts (By similarity. Acts as a negative regulator of antigen cross-presentation in myeloid dendritic cells (By similarity. In the context of tumorigenesis, negative regulation of antigen cross-presentation limits the anti-tumor response by reducing efficiency of tumor-antigen cross-presentation (By similarity.
基因ID	54915
基因名	YTHDF1
Swiss	Q9BYJ9
别名	YTHDF1;C20orf21

产品验证



Western blot analysis of YTHDF1 expressed in K-562, Mouse testis, Rat thymus using YTHDF1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)