

Histone H3 Rabbit mAb

货号: B11632

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	WB: Homo sapiens , Mus musculus , Sus scrofa
应用	WB IHC CHIP
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 ChIP: 1:50 - 1:200
理论分子量	15kDa
实测分子量	17KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,Raji,NIH/3T3,C6,Mouse liver,Rat spleen,Rat lung
细胞定位	extracellular exosome,extracellular region,nucleoplasm,nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide of human Histone H3 (NP_003520.1).
序列	RTKQT

靶点信息

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosom e, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucle osomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.
基因ID	8350
基因名	
Swiss	P68431
别名	H3/A; H3FA

产品验证



Western blot analysis of Histone H3 expressed in Mouse liver,Rat spleen,HeLa using Histone H3 Rabbi t mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 3 0ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure tim e: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn