

Leptin Rabbit pAb

货号: B11816

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Human colon cancer cell , Homo sapiens IHC: Human colon cancer cell Electrochemical Sensing: Rattus norvegicus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	18kDa
实测分子量	15KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,BxPC-3,RD,Mouse kidney,Rat testis
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Leptin (NP_000221. 1).	
序列	MHWGTLCGFLWLWPYLFYVQAVPIQKVQDDTKTLIKTIVTRINDISHTQSVSSKQKVTGLDFIPGLHPILTLSKMDQTLAVY QQILTSMPSRNVIQISND	

靶点信息

研究背景	This gene encodes a protein that is secreted by white adipocytes, and which plays a major role in the regulation of body weight. This protein, which acts through the leptin receptor, functions as part of a signalin g pathway that can inhibit food intake and/or regulate energy expenditure to maintain constancy of the a dipose mass. This protein also has several endocrine functions, and is involved in the regulation of immune and inflammatory responses, hematopoiesis, angiogenesis and wound healing. Mutations in this gene a nd/or its regulatory regions cause severe obesity, and morbid obesity with hypogonadism. This gene has also been linked to type 2 diabetes mellitus development.
基因ID	3952
基因名	LEP
Swiss	P41159
别名	LEP;LEPD;OB;OBS;leptin

产品验证



Western blot analysis of Leptin expressed in Mouse kidney,Rat testis,HepG2 using Leptin Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lan e. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn