

# **CAMKK2 Rabbit pAb**

货号: **B11820** 

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Rabbit cerebral microvascular endothelial cells , Homo sapiens , Mus musculus , Macaca mulatta
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	54kDa/58kDa/59kDa/61kDa/64kDa
实测分子量	68KDa/70KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,SH-SY5Y,LNCaP
细胞定位	Cell projection,Cytoplasm,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-100 of human CAMKK 2 (NP_006540.3).	
序列	MSSCVSSQPSSNRAAPQDELGGRGSSSSESQKPCEALRGLSSLSIHLGMESFIVVTECEPGCAVDLGLARDRPLEADGQE VPLDTSGSQARPHLSGRKLS	

靶点信息

研究背景	The product of this gene belongs to the Serine/Threonine protein kinase family, and to the Ca(2+)/calmod ulin-dependent protein kinase subfamily. The major isoform of this gene plays a role in the calcium/calmo dulin-dependent (CaM) kinase cascade by phosphorylating the downstream kinases CaMK1 and CaMK4. P rotein products of this gene also phosphorylate AMP-activated protein kinase (AMPK). This gene has its str ongest expression in the brain and influences signalling cascades involved with learning and memory, ne uronal differentiation and migration, neurite outgrowth, and synapse formation. Alternative splicing result s in multiple transcript variants encoding distinct isoforms. The identified isoforms differ in their ability to undergo autophosphorylation and to phosphorylate downstream kinases.
基因 <b>ID</b>	10645
基因名	CAMKK2
Swiss	Q96RR4
别名	CAMKK2;CAMKKB

## 产品验证



Western blot analysis of CAMKK2 expressed in HeLa,SH-SY5Y,LNCaP using CAMKK2 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

# 实验步骤

访问官网浏览详情: www.ablybio.cn