



Smad2 Rabbit pAb

货号: **B11841**

产品信息

| 反应 | Human,Mouse,Rat |
|---------|---|
| 12,127. | numan, Mouse, Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: U87MG , Toxoplasma gondii , Rattus norvegicus , Homo sapiens |
| 应用 | WB IHC IF/ICC IP |
| 推荐浓度 | WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:20 - 1:50 IP: 1:20 - 1:50 |
| 理论分子量 | 48kDa/52kDa |
| 实测分子量 | 50kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa |
| 细胞定位 | Cytoplasm,Nucleus |
| 纯化 | Affinity purification |

抗原信息

| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 175-270 of human Sma d2 (NP_005892.1). | |
|------|--|--|
| 序列 | LPPVLVPRHTEILTELPPLDDYTHSIPENTNFPAGIEPQSNYIPETPPPGYISEDGETSDQQLNQSMDTGSPAELSPTTLSPVN HSLDLQPVTYSE | |

靶点信息

| 研究背景 | The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD prot eins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cell ular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TG F-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In r esponse to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it bin ds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Altern atively spliced transcript variants have been observed for this gene. |
|--------------|--|
| 基因 ID | 4087 |
| 基因名 | SMAD2 |
| Swiss | Q15796 |
| 别名 | JV18;JV18-1;MADH2;MADR2;hMAD-2;hSMAD2;Smad2;SMAD2 |

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn