

# Histone H2AX Rabbit pAb

## 货号**: B12176**

### 产品信息

反应	Human,Mouse,Rat,Other (Wide Range)
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: U87MG , Homo sapiens Western Blot: Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100
理论分子量	15kDa
实测分子量	15kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,MCF7,HL-60,Rat heart
细胞定位	Chromosome,Nucleus
纯化	Affinity purification

### 抗原信息

抗原信息	A synthetic phosphorylated peptide around S2 of human Histone H2AX (NP_002096.1).	
序列	MSGRG	

#### 靶点信息

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octa mer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The l inker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chr omatin into higher order structures. This gene encodes a replication-independent histone that is a memb er of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop t ermination motif, and the polyA addition motif.
基因ID	3014
基因名	H2AFX
Swiss	P16104
别名	H2A.X;H2A/X;H2AX;Histone H2AX;H2AFX;histone H2AX;

#### 产品验证



Western blot analysis of Histone H2AX expressed in Rat heart,293T,MCF7 using Histone H2AX Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30 ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure tim e: 120s.

#### 实验步骤

访问官网浏览详情: www.ablybio.cn