

ISG15 Rabbit pAb

货号: **B12262**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Dragon Li cats , Bos taurus
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200
理论分子量	17kDa
实测分子量	15KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	RD,Jurkat
细胞定位	Cytoplasm,Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-165 of human ISG15 (NP_005092.1).	
序列	MGWDLTVKMLAGNEFQVSLSSSMSVSELKAQITQKIGVHAFQQRLAVHPSGVALQDRVPLASQGLGPGSTVLLVVDKC DEPLSILVRNNKGRSSTYEVRLTQTVAHLKQQVSGLEGVQDDLFWLTFEGKPLEDQLPLGEYGLKPLSTVFMNLRLRGGG TEPGGRS	

靶点信息

研究背景 The protein encoded by this gene is a ubiquitin-like protein that is conjugated to intracellular target pr ns upon activation by interferon-alpha and interferon-beta. Several functions have been ascribed to th ncoded protein, including chemotactic activity towards neutrophils, direction of ligated target proteins intermediate filaments, cell-to-cell signaling, and antiviral activity during viral infections. While conjug s of this protein have been found to be noncovalently attached to intermediate filaments, this protein ometimes secreted.
基因 ID 9636
基因名 ISG15
Swiss P05161
別名 ISG15;G1P2;IFI15;IMD38;IP17;UCRP;hUCRP

产品验证



Western blot analysis of ISG15 expressed in RD,Jurkat using ISG15 Rabbit pAb at 1:1000. Secondary antib ody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn